

The Outcome of Intralesional Steroid Injection for Ganglion Cyst compared to surgery

Khwaja H Salim*,¹ Zobaer Hassan Chowdhury,² Shams E Tabriz,³ Md. Humayun Kabir.⁴

ABSTRACT: Background: Ganglia are the most common benign cystic swellings found over both the dorsal and volar aspects of the wrist. In spite of technical advancement, both operative and non-operative interventions achieve more or less similar results. Complete evacuation of gelatinous fluid followed by intralesional injection of Triamcinolone gives encouraging results. This study was done to explore the outcome of intralesional steroid injection compared to surgery for the management of ganglion cyst. **Materials and Methods:** A prospective study was conducted on patients with simple ganglion cysts on the wrist. Total of 100 patients underwent this study. Out of this, 45 patients underwent aspiration of the cyst fluid followed by intracystic instillation of triamcinolone, and 55 patients underwent surgical excision of wrist ganglia. Two years follow up was done for evaluation of outcome. **Result:** Most ganglia of wrist occurred in the extensor aspect. Complications noted among the surgically excised group were post operative pain and restricted mobility of wrist with a recurrence rate of 21.5%. Instillation of triamcinolone into the ganglion yielded early resolution with a low recurrence rate of 8.4%. **Conclusion:** Recurrence is the most common complication of management of ganglion cyst & its incidence is less following intralesional triamcinolone injection compared to surgical excision.

Keywords: Ganglion cyst, triamcinolone, intralesional injection

(The Planet 2018; 2(2): 27-32)

INTRODUCTION:

Wrist ganglia are the benign soft tissue tumors most commonly encountered on the dorsal aspect of the wrist and may communicate with the joint via a pedicle. Ganglia found on the volar aspect of the wrist attach via a pedicle to the

radioscaphoid, scapholunate, scaphotrapezial joint, or the metacarpotrapezial joint.¹ They are cystic structures measuring 1-2 cm in diameter, feel much like a firm nodular swelling that is well tethered in place by its attachment to the underlying joint capsule or tendon

1. Assistant Professor, (Pediatric Surgery), Shaheed Tazuddin Ahmad Medical College, Gazipur.

2. Assistant Professor (Pediatric Surgery), Sheikh Sayera Khatun Medical College, Gopalganj.

3. Associate Professor (Anatomy), Sheikh Sayera Khatun Medical College, Gopalganj.

4. Associate Professor (Ortho Surgery), Sheikh Sayera Khatun Medical College, Gopalganj.

* For correspondence- Dr. Khwaja H Salim, Assistant Professor, (Pediatric Surgery), Shaheed Tazuddin Ahmad Medical College, Gazipur.

sheath. Symptoms include pain in the wrist, which may radiate to the arm, and may be exacerbated by wrist movement, decreased range of motion, and reduced grip strength. Volar ganglia can also cause paresthesias from compression of the ulnar or median nerves or their branches.²

Surgery is the mainstay of therapeutic option employed for managing ganglia and involves excision of the entire ganglion complex, including cyst, pedicle, and a cuff in the adjacent joint capsule. Recurrence often occurs and is attributed to the remnants of the tortuous duct system located near the joint capsule due to inadequate dissection.

We describe a simple double puncture method of instillation of triamcinolone as a safe and more effective method for management of wrist ganglion.

MATERIAL & METHOD:

A prospective study was conducted on patients attending surgical outpatient department at Shaheed Tazuddin Ahmad Medical College Hospital (STAMCH) for simple ganglion cysts on the wrist from March 2017 to June 2018. A total of 105 patients were recruited in this study. Out of this, 52 patients underwent the aspiration of the cyst fluid followed by intralesional injection of triamcinolone, and 53 patients underwent surgical excision of wrist ganglia. We included all the cases of wrist ganglia both on extensor as well as flexor aspect of the wrist. More extensive ganglia like compound palmar ganglia were excluded. Cysts were examined for tenseness and cystic consistency by clinical examination.

One sterile wide bore IV cannula of size 18 gauge and another 23 gauge hypodermic needles were inserted into the cyst wall facing each other. The gelatinous content was drained through the cannula by a combination of negative pressure with 10cc disposable syringe & gentle compression of the cyst. After complete evacuation of cyst contents, the cannula was withdrawn and the puncture site was sealed with a swab. Triamcinolone acetonide (Trialon) 40mg/ml was injected into the cyst cavity via previously placed size 23G hypodermic needle, thus again refilling the cyst cavity to attain similar size. Overstretching of the cyst wall was avoided. This needle was withdrawn and the puncture site was sealed with spirit swab. Compression was applied over the injected cyst. Patients were followed up every month for six months. Maximum follow up period was up to two years.

RESULTS:

Out of 105 patients, 74 were females and 31 were males showing female preponderance. Age of patients ranged from 8 to 37 years with the highest incidence noted in the age group between 15 to 25 years (44/105, 41.9%). Ganglia on extensor aspect were more common than on volar surface (91/105, 86.6%) in our study. In the triamcinolone group, there was mild pain locally for 48 hours which responded well to oral analgesics. None of them had any restricted wrist mobility, joint instability, or hypopigmentation of overlying skin. There was recurrence in 4 patients out of 52 (7.6%) within 6 months of the procedure, which responded to repetition of local

triamcinolone injection. There were no steroid induced side effects. 8 out of 53 patients of surgical excision cohort had recurrence (15%). All these cases had post operative pain, lasting for four to five days and required analgesics. Wrist movements remained restricted for three to four weeks.

DISCUSSION:

Ganglia are localized, tense but often painless cystic swellings containing clear gelatinous fluid. They often communicate with and are always adjacent to a tendon sheath or the capsule of a joint. Their origin is uncertain, but they are probably caused by myxoid degeneration of fibrous tissue of capsule, ligaments and retinaculae. It consists of an outer fibrous coat and an inner synovial lining.¹ These occur singly but may be unilocular or multilocular. Ganglia are more common in women and often occur between second and third decades (70%).

Ganglia are frequently located in the hand and wrist and less often in the ankle, foot and knee. In the hand, 60-70% of all ganglia occur over scapholunate ligament. Nearly 20% of them present over volar aspect of wrist crease between flexor carpi radialis and abductor pollicis longus at the scaphotrapezoid joint.^{2,3} There may be scapholunate diastasis that may persist even after excision.

Clinically, they present as slowly growing localized swellings with mild aches. They range from smooth, spherical and semi fluctuant to firm in consistency, with positive transillumination. They become more prominent on flexing the wrist. The

extent of cyst and direction of pedicle can be well made out on compressing the cyst.⁴

Ganglia of the wrist may have a variable appearance on sonography like the presence of lobulations, well-defined margins, internal echoes, and septations. Small ganglion cysts often appear hypoechoic without posterior acoustic enhancement, not appearing as simple cysts.⁵

Various management modalities are available. Simple technique of aspiration with a short beveled wide bore needle has been tried, with about 60% recurrence within 3 months.⁶ Repeated aspirations gave 85% success rate. Many patients gave up the procedure during follow-ups. Clay and Clement performed a radical excision of wrist ganglia by excising attachments up to scapholunate ligament. But this technique was associated with persistence of discomfort and recurrence with joint instability.⁷ Regardless of whether the ganglion is removed via open surgery or arthroscopic excision when a stalk is visualized, it should be completely removed, or the cyst should be removed at the origin to reduce the risk of recurrence. A recurrence rate of 13-40% has been observed following removal of entire cysts.⁸

A volar ganglion cyst is much more likely to recur than a dorsal ganglion cyst and care must be taken to protect the radial and ulnar neurovascular bundles because radial artery laceration is the most common vascular complication in volar carpal excision of ganglia. There was evidence of nerve damage to the superficial branch of the radial nerve and to the palmar

cutaneous branch of the median nerve.⁸ In open and arthroscopic procedures, wrist stiffness is the most common postoperative complication, along with infection, decreased motion, and ligament instability in open especially of the scapholunate ligament.^{9,10,11} Arthroscopic resection of volar ganglion of wrist has been done with comparable results. In some, cyst wall was not removed and wide stalk has been left behind. The average operative time for such case has been 55 minutes.¹² Even then, few recurrences have occurred. Rizzo reported stiffness in 25% of patients and required up to 8 weeks of occupational therapy to regain maximal function.¹³

Local steroid infiltration in ganglion has shown satisfactory results in adults as well as in children.^{14,15,16} Hyaluronidase also has been used to improve liquefaction in the cysts.^{17,18} Steroids probably arrest the secretions from mesenchymal cells at the cyst wall. In our series, evacuation of gelatinous fluid resulted in complete emptying of cyst cavity. Instilled triamcinolone, because of its close contact with mucin secreting epithelium, leads to a cessation of secretion of gelatinous fluid.

CONCLUSION:

We have evaluated a simple, noninvasive, safe and effective method of treatment for this common condition. This double puncture procedure is scarless with a lower rate of recurrence and therefore easily acceptable to the patients. The absence of side effects such as wrist stiffness, joint instability or pain is an added advantage.

REFERENCES:

1. Plate AM, Lee SJ, Steiner G, Posner MA. Tumor like lesions and benign tumors of the hand and wrist. *J Am Acad Orthop Surg.* 2003;11:129-41.
2. Thornburg LE. Ganglions lesions of the hand and wrist. *J Am Acad Orthop Surg.* 1999;7: 231-8.
3. Angelides AC, Wallace PF. The dorsal ganglion of the wrist: Its pathogenesis, gross and microscopic anatomy, and surgical treatment. *J Hand Surg.* 1976;1:228-35.
4. Nahra ME, Bucchieri JS. Ganglion cysts and other tumor related conditions of the hand and wrist. *Hand Clin.* 2004;20:249-60.
5. Wang G, Jacobson JA, Feng FY, Girish G, Caoili EM, Brandon C. Sonography of wrist ganglion cysts: Variable and noncystic appearances. *J Ultrasound Med.* 2007;26:1323-8.
6. Zubowicz VN, Ischii CH. Management of ganglion cysts of the hand by simple aspiration. *J Hand Surg.* 1987;12:618-20.
7. Clay NR, Clement DA. The treatment of dorsal wrist ganglia by radical excision. *J Hand Surg.* 1988;13:187-91.
8. Edwards SG, Joansen JA. Prospective outcomes and associations of wrist ganglion cysts resected arthroscopically. *J Hand Surg Am.* 2009;34: 395-400.
9. Rocchi L, Canal A, Fanfani F, Catalano F. Articular ganglia of the volar aspect of the wrist: Arthroscopic resection compared with open excision: A prospective randomised study. *Scand J Plast Reconstr Surg Hand Surg.* 2008;42:253-9.
10. Mehdian H, McKee MD. Scapholunate instability following dorsal wrist ganglion excision. *Iowa Orthop J.* 2005;25:203-6.

11. Duncan KH, Lewis RC. Scapholunate instability following ganglion cyst excision: A case report. *Clin Orthop Relat Res.* 1988;228:250-3.
12. Nishikawa S, Toh S, Miura K, Arai K, Irie T. Arthroscopic diagnosis and treatment of dorsal wrist ganglion. *J Hand Surg.* 2001;26:547-9.
13. Rizzo M, Berger RA, Steinmann SP, Bishop AT. Arthroscopic resection in the management of dorsal wrist ganglions: Results with a minimum 2-year follow-up period. *J Hand Surg Am.* 2004;29:59-62.
14. Neidoff M, Davis TRC, Clay NR. Conservative management of wrist ganglia: Aspiration versus steroid infiltration. *J Hand Surg.* 1997;22:636-7.
15. Colberg RE, Sánchez CF, Lugo-Vicente H. Aspiration and triamcinolone acetonide injection of wrist synovial cysts in children. *J Pediatr Surg.* 2008;43:2087-90.
16. Paramhans D, Nayak D, Mathur RK, Kushwah K. Double Dart Technique of Instillation of Triamcinolone in Ganglion Over the Wrist. *J Cutan Aesthet Surg.* 2010 Jan-Apr; 3(1): 29-31
17. Jagers M, Akkerhuis P, van Der Heijden M, Brink PRG. Hyaluronidase versus surgical excision of ganglia: a prospective, randomized clinical trial. *J Hand Surg.* 2002;27:256-8.
18. Paul AS, Sochart DH. Improving the results of ganglion aspiration by the use of hyaluronidase. *J Hand Surg.* 1997;22:219-21.