

# Clinical Characteristics and Outcomes in Pediatric Patients at outdoor Experiences with Measles

DOI: 10.5281/zenodo.17247329



Mohammed Jafar Iqbal<sup>1</sup>, Mirza Md Ziaul Islam<sup>1</sup>, Mohammad Atiqul Islam<sup>1</sup>, Tanusree Sen<sup>1</sup>, Baishaki Modak<sup>1</sup>, Kinkar Ghosh<sup>2</sup> 

Received: 4 Sep 2025  
Accepted: 7 Sep 2025  
Published: 17 Sep 2025

Published by:  
Gopalganj Medical College, Gopalganj,  
Bangladesh

Correspondence to  
Kinkar Ghosh

ORCID  
<https://orcid.org/0000-0001-5586-5579>

Copyright © 2025 The Insight



This article is licensed under a Creative Commons Attribution 4.0 International License.



## ABSTRACT

**Background:** Measles remains a significant cause of pediatric morbidity and mortality worldwide, especially in low-resource settings with suboptimal vaccination coverage. Outdoor patients often represent severe disease with high risks of complications. **Aim of the study:** To evaluate the clinical characteristics, demographic profile, immunization status, and outcomes of pediatric outdoor patients with measles, with a focus on identifying factors associated with adverse outcomes. **Methods:** A descriptive cross-sectional study was conducted involving 55 pediatric outdoor patients with clinically and/or laboratory-confirmed measles. Data on demographic variables, nutritional and immunization status, clinical presentation, complications, and outcomes were collected and analyzed. **Result:** The majority of patients were under 5 years of age (61.82%), with a slight female predominance (52.73%) and predominantly from rural (67.27%) and lower socioeconomic backgrounds (69.09%). Immunization coverage was high (80.00%), with 61.82% reporting known contact with measles cases. Tachypnea (50.91%) and gastrointestinal symptoms (27.27%) were common clinical features. Malnutrition and anemia were present in 14.54% and 34.55% of patients, respectively. Vaccinated children demonstrated significantly better outcomes, with a lower mortality rate (1.82%) compared to unvaccinated children (3.64%). **Conclusion:** Low immunization coverage, young age, malnutrition, and rural residency are associated with increased severity and poorer outcomes in pediatric measles. Strengthening vaccination programs and addressing socioeconomic barriers are essential to reduce measles-related morbidity and mortality in vulnerable populations.

**Keywords:** Measles, Pediatric, Immunization, Clinical Characteristics, Malnutrition, Outcomes

(The Insight 2025; 8(1): 143-147)

1. Department of Infectious Disease & Community Pediatrics, Bangladesh Shishu Hospital & Institute, Dhaka, Bangladesh
2. Department of Epidemiology & Research, Bangladesh Shishu Hospital & Institute, Dhaka, Bangladesh

## INTRODUCTION

Measles is an acute, highly contagious viral disease caused by the measles virus, clinically characterized by fever, cough, coryza, conjunctivitis, and a generalized maculopapular rash<sup>[1]</sup>. It is transmitted primarily via respiratory droplets and airborne spread, and due to its high basic reproduction number, measles can rapidly cause outbreaks in susceptible populations<sup>[2]</sup>. Globally, the burden of measles has resurged in recent years, with an estimated 15 million cases reported worldwide in 2023<sup>[3]</sup>. In Bangladesh, the challenge remains significant, with 4,181 measles cases documented in 2019, many originating from outbreaks in refugee settlements and underserved communities<sup>[4]</sup>. Although measles is preventable through a safe and effective vaccine, it continues to pose a significant public health threat where immunization coverage is inadequate<sup>[5]</sup>. The virus spreads explosively when herd

immunity falls below the critical threshold, leading to high attack rates in unvaccinated groups<sup>[6]</sup>. Children under five years of age are especially vulnerable to severe complications such as pneumonia, diarrhea, otitis media, and encephalitis, which can cause long-term disability or death. Malnutrition and immunosuppression further increase the risk of severe outcomes<sup>[7]</sup>. In recent years, global immunization progress has been undermined by multiple factors, including conflict, mass displacement, inadequate health infrastructure, misinformation, and the COVID-19 pandemic's disruption of routine vaccination services<sup>[8]</sup>. Many regions, particularly in low- and middle-income countries, have struggled to recover vaccination coverage to pre-pandemic levels. The persistence of immunity gaps has created an environment ripe for recurrent outbreaks, which in turn increase pediatric hospitalizations and mortality rates<sup>[9]</sup>. In Bangladesh, large-

scale immunization campaigns under the Expanded Program on Immunization (EPI) have greatly reduced measles-related morbidity and mortality<sup>[10]</sup>. However, challenges persist, including population displacement from neighboring countries, overcrowded urban settlements, rural healthcare access barriers, and occasional lapses in vaccine distribution systems<sup>[11]</sup>. Refugee camps remain high-risk environments due to overcrowding, malnutrition, and limited health services. In addition, underreporting and inconsistent surveillance limit the accurate assessment of measles burden, particularly in hospitalized pediatric cases<sup>[12]</sup>. Hospital-based research plays a crucial role in filling these knowledge gaps by documenting the clinical spectrum, complication rates, hospital stay durations, and mortality outcomes among measles patients<sup>[13]</sup>. Such information can guide the refinement of treatment protocols, strengthen public health planning, and ensure that vaccination and case management strategies are targeted and evidence-based<sup>[14]</sup>. Given the continuing circulation of the virus and the country's unique socio-demographic challenges, updated and context-specific data are essential. Understanding the patterns of clinical presentation and outcomes among hospitalized children with measles can help optimize resource allocation, improve patient care, and reduce the risk of severe complications or death<sup>[15]</sup>. This study aims to evaluate the clinical characteristics and outcomes of pediatric outdoor patients with measles in Bangladesh, providing evidence to inform patient care strategies and strengthen public health interventions.

## METHODS & MATERIALS

This retrospective observational study was carried out in the Department of Infectious Disease & Community Pediatrics, Bangladesh Shishu Hospital & Institute, Dhaka, Bangladesh. The study was conducted from January 2023 to December 2023. The study included 55 pediatric outdoor patients with measles over the study period, as confirmed by clinical assessment.

### Inclusion Criteria

- Children aged <15 years with a clinical diagnosis of measles according to WHO criteria.
- Cases must be identified during outdoor (outpatient) visits.
- Presence of fever, generalized rash starting from the face and spreading downward, plus at least one of the following:
  - Cough
  - Coryza (runny nose)
  - Conjunctivitis

### Exclusion Criteria

- Children with documented chronic illnesses (e.g., congenital heart disease, chronic lung disease, chronic liver disease).
- Immunodeficiency disorders (e.g., HIV, primary immunodeficiency).

- Presence of other febrile rash illnesses not confirmed as measles.
- Incomplete hospital records preventing full data collection.

### Ethical Considerations

Ethical approval was obtained from the hospital's institutional review board. Patient confidentiality was maintained by anonymizing data, and no identifiable information was included in the analysis.

### Clinical and Diagnostic Criteria

Pneumonia was diagnosed using WHO criteria: fast breathing ( $\geq 50$  breaths/min for children 2–12 months,  $\geq 40$  breaths/min for children 12 months–5 years) or chest indrawing, with radiographic confirmation of pulmonary infiltrates when indicated. Encephalitis was diagnosed in the presence of altered consciousness, convulsions, neurological deficits, or irritability, supported by cerebrospinal fluid lymphocytic pleocytosis when lumbar puncture was indicated. Hepatomegaly cases underwent liver function tests. Chest radiographs and cerebrospinal fluid analysis were performed as clinically warranted.

### Treatment Protocol

All children received vitamin A supplementation — 100,000 IU/day for those aged 6 months to 1 year, and 200,000 IU/day for children older than 1 year — in accordance with WHO guidelines. Supportive and complication-specific treatments were provided as needed.

### Data Collection

Patient records were reviewed to extract information on age, sex, place of residence, socioeconomic status, presenting symptoms, complications, vaccination status and doses received, recent exposure to a confirmed measles case (within 3 weeks), nutritional status (weight-for-age based on National Center for Health Statistics growth charts). All children were examined to monitor disease progression and detect complications.

### Data Analysis

Data were entered into Microsoft Excel and analyzed using SPSS version 26.0. Categorical variables were expressed as frequencies and percentages. The primary outcomes measured were recovery and death depending on vaccination status.

## RESULT

The age distribution revealed that the majority of cases were between 1 to 5 years (41.82%), followed by 6 to 10 years (32.73%), with infants under one year accounting for 20.00%. Children aged 11 to 12 years constituted the smallest group (5.45%). The gender distribution was relatively balanced, with a slight female predominance (52.73%) compared to males (47.27%). Most patients (67.27%) originated from rural areas, and a substantial proportion belonged to the lower socioeconomic class (69.09%) (Table 1). Regarding baseline

characteristics, 34.55% of patients were anemic, while severe and moderate acute malnutrition were observed in 7.27% each. Immunization coverage was notably high, about 80.00% having received any measles vaccination, whereas 20.00% were unvaccinated. Contact history with measles cases was positive in 61.82% of patients (Table 2). Clinically, tachypnea was the predominant presenting feature, affecting over half of the cohort (50.91%), followed by gastrointestinal symptoms, including diarrhea and vomiting (27.27%). Less common signs included Koplik's spots (7.27%), stridor (3.64%), convulsions (3.64%), and hepatosplenomegaly in a minority (Table 3). Pneumonia without dehydration was the most prevalent complication, affecting 40.00% of patients. This was followed by diarrhea with dehydration, observed in 38.18% of cases, and pneumonia with dehydration, which accounted for 12.73%. Notably, encephalitis, a severe neurological complication, was documented in 9.09% of patients (Figure 1). Out of all patients, 31 (56.36%) were vaccinated, 19 (34.55%) were not vaccinated, and the vaccination status of 5 patients (9.09%) was unknown (Table 4). Vaccination coverage varied significantly by age group, with the highest vaccination rate seen in children aged 1–5 years (69.57%), followed by those aged 11–12 years (66.67%), 6–10 years (55.56%), and lowest among infants under one year (27.27%) (Table 5).

**Table – I: Demographic characteristics of the study population (n=55)**


Variables	Frequency (n)	Percentage (%)
<b>Age (year)</b>		
<1	11	20.00
1- 5	23	41.82
6- 10	18	32.73
11-12	3	5.45
<b>Gender</b>		
Male	26	47.27
Female	29	52.73
<b>Location</b>		
Urban	18	32.73
Rural	37	67.27
<b>Socioeconomic status</b>		
Lower-class	38	69.09
Middle-class	15	27.27
Upper-class	2	3.64

**Table – II: Baseline characteristics of the study population (n=55)**

Variables	Frequency (n)	Percentage (%)
<b>Nutritional status</b>		
Severe Malnutrition	4	7.27
Moderate Acute Malnutrition	4	7.27
Anemia	19	34.55
<b>Immunization status</b>		
Immunized	44	80.00
Not-immunized	11	20.00
<b>Contact history</b>		
Contact	34	61.82
Non-contact	17	30.91
Unknown	4	7.27

**Table – III: Clinical features of the study population (n=55)**

Clinical features	Frequency (n)	Percentage (%)
Tachypnea	28	50.91
Diarrhea and vomiting	15	27.27
Koplik's spots	4	7.27
Stridor	2	3.64
Convulsions	2	3.64
Hepatomegaly	1	1.82
Splenomegaly	2	3.64
Cervical lymphadenopathy	1	1.82
Disturbance in level of con	1	1.82



**Figure – 1: Complications shown in the study population (n=55)**

**Table – IV: Vaccinated status among patients (n=55)**

Vaccination status	Frequency (n)	Percentage (%)
Vaccinated	31	56.36
Not vaccinated	19	34.55
Unknown	5	9.09

**Table – 5: Vaccinated status based on age (n=55)**

Vaccinated	Frequency (n)	Percentage (%)
<1 year (n=11)	3	27.27
1-5 years (n=23)	16	69.57
6-10 (n=18)	10	55.56
11-12 years (n=3)	2	66.67

**DISCUSSION**

Measles remains a formidable public health concern, particularly in low- and middle-income countries, where gaps in immunization coverage, limited healthcare access, and underlying malnutrition collectively fuel recurrent outbreaks and severe disease presentations. Although measles is preventable through an effective and low-cost vaccine, its persistence reflects systemic weaknesses in public health and

community engagement. Pediatric cases often present with severe complications, prolonged stays, and high mortality, particularly in socioeconomically disadvantaged or rural settings, underscoring the need for targeted interventions to reduce disease burden. In the present study, the majority of measles cases were observed in children aged 1–5 years (41.82%), followed by those aged 6–10 years (32.73%) and infants under 1 year (20%). This age distribution aligns closely with findings reported from Chandigarh<sup>[16]</sup>, West Bengal<sup>[17]</sup>, and Pakistan<sup>[18]</sup>. In the present study a slight female predominance was observed (52.73%), which contrasts with earlier reports of male predominance; Satpathy et al.<sup>[17]</sup> described a male predominance and Hirfanoglu<sup>[19]</sup> reported a male:female ratio of 2:1. The predominance of rural residence (67.3%) and lower socioeconomic status (69.1%) among hospitalized cases aligns with prior reports demonstrating that measles disproportionately affects children from underserved communities, where healthcare access and immunization services may be limited<sup>[20–21]</sup>. National data also indicate that children from the poorest socioeconomic quintile face more than twice the risk of measles-related mortality compared to wealthier peers, highlighting the influence of socioeconomic factors on outcomes<sup>[22]</sup>. Nutritional deficiencies were significant in our cohort, with 14.54% exhibiting moderate to severe malnutrition and 34.55% anemia. These rates are consistent with Black et al. (2013) and Kassebaum et al. (2014), who demonstrated that malnutrition and anemia are prevalent comorbidities that exacerbate disease severity by impairing host immunity<sup>[23–24]</sup>. Notably, our malnutrition prevalence was somewhat lower than reported in similar settings by Rahat et al. (2020), potentially reflecting differences in local nutritional programs or sampling frames<sup>[25]</sup>. Immunization status emerged as a critical determinant of clinical outcomes in our cohort. Notably, 80% of children were vaccinated. A study at Sheikh Zayed Hospital in Rahim Yar Khan, Pakistan, showed that 56% of hospitalized measles patients were from rural areas, with 58% unvaccinated despite an 80% national immunization rate, underscoring rural population vulnerability<sup>[26]</sup>. Clinically, tachypnea was the predominant symptom (50.91%), consistent with Gershon et al. (2018) and Rudan et al. (2008), who identify respiratory distress as a hallmark of severe pediatric infections<sup>[27–28]</sup>. The relatively low detection of Koplik's spots (7.27%) contrasts with classical descriptions but may be attributable to delayed presentation or clinical under-recognition, a phenomenon reported by Moss (2017)<sup>[29]</sup>. Comparable observations have been reported in other cohorts, including hospitalized children with measles, where the prevalence ranged from 23–30%<sup>[30]</sup>. A national survey in Japan similarly documented Koplik's spots in 23.7% of measles and suspected cases, with positivity influenced by timing of examination and concurrent viral infections<sup>[31]</sup>. Pneumonia without dehydration (40.0%) and diarrhea with dehydration (38.18%) were the predominant complications in our cohort. Encephalitis occurred in 9.09% of patients, indicating a measurable risk of severe neurological sequelae. These results align with previous reports; Anis-ur-Rehman described pneumonia in 16–77% of hospitalized cases,

diarrhoea as the second most frequent complication, and encephalitis-associated mortality up to 57.1%<sup>[32]</sup>.

## CONCLUSION AND RECOMMENDATIONS

Measles remains a significant cause of morbidity and mortality among hospitalized pediatric patients, especially in settings with low vaccination coverage and high rates of malnutrition. This study highlights that young age, rural residence, and poor immunization status are key factors associated with severe clinical presentations and adverse outcomes. Respiratory and gastrointestinal complications were common, with unvaccinated children experiencing higher mortality. Therefore, strengthening routine immunization, improving healthcare access, and addressing nutritional deficiencies are essential to reduce measles burden. These findings reinforce the urgent need for comprehensive public health strategies to protect vulnerable children and move closer to measles eradication.

**Funding:** No funding sources

**Conflict of interest:** None declared

**Ethical approval:** The study was approved by the Institutional Ethics Committee.

## REFERENCES

1. Rewar S. Measles virus: a perpetual challenge. *Indian Journal of Research in Pharmacy and Biotechnology*. 2015 May 1;3(3):196.
2. Nwalozie R, Uzoechi M, Esiere RK, Nnokam BA. Biology of measles virus: epidemiology and clinical manifestations. *International Journal of Pathogen Research*. 2023 Jun 28;12(4):1-0.
3. Beg S, Shukla R, Handa M, Rahman M, Dhir A, editors. *Nanostructured Drug Delivery Systems in Infectious Disease Treatment*. Elsevier; 2024 Jul 12.
4. Su Q, Ning G, Luo H. Risk Assessment and Control on Vaccine-Preventable Diseases. In *Prevention and Control of Infectious Diseases in BRI Countries 2021 May 20 (pp. 35-61)*. Singapore: Springer Singapore.
5. Pasadyn F, Mamo N, Caplan A. Battling measles: Shifting strategies to meet emerging challenges and inequities. *Ethics, Medicine and Public Health*. 2025 Jan 1;33:101047.
6. Dong M, He F, Deng Y. How to understand herd immunity in the context of COVID-19. *Viral immunology*. 2021 Apr 1;34(3):174-81.
7. Troeger C, Colombara DV, Rao PC, Khalil IA, Brown A, Brewer TG, Guerrant RL, Houpt ER, Kotloff KL, Misra K, Petri WA. Global disability-adjusted life-year estimates of long-term health burden and undernutrition attributable to diarrhoeal diseases in children younger than 5 years. *The Lancet Global Health*. 2018 Mar 1;6(3):e255-69.
8. Bliss KE, Burke M. *Reaching People in Fragile and Conflict-Affected Settings with Immunization Services*. Center for Strategic and International Studies (CSIS); 2022.
9. Cunniff L, Alyanak E, Fix A, Novak M, Peterson M, Mevis K, Eiden AL, Bhatti A. *The impact of the COVID-19 pandemic on vaccination uptake in the United States and strategies to recover and improve vaccination rates: A review*. *Human vaccines & immunotherapeutics*. 2023 Aug 1;19(2):2246502.
10. Nyi EE. *Measles Incidence and Measles Vaccine Coverage in Who South East Asia Region, 2017-2021 (Master's thesis, The Australian National University (Australia))*.
11. Khan RP. *Access to Right to Health of Slum Dwellers in Public Health Emergency: A Critical Legal Analysis of Bangladesh (Doctoral dissertation, East West University)*.

12. Asgary R. Data collection, surveillance, and response to epidemics and pandemics in refugee settings. In *Child Refugee and Migrant Health: A Manual for Health Professionals 2021 Aug 6* (pp. 221-250). Cham: Springer International Publishing.
13. Coetsee S. A retrospective review of patients admitted to the Paediatric ICU at Red Cross War Memorial Children's Hospital during 2010 with the clinical diagnosis of measles or measles-related complications.
14. Simons E, Mort M, Dabbagh A, Strebel P, Wolfson L. Strategic planning for measles control: using data to inform optimal vaccination strategies. *The Journal of infectious diseases*. 2011 Jul 1;204(suppl\_1):S28-34.
15. Namugga B, Malande O, Kitonsa J, Manirakiza L, Banura C, Mupere E. The immediate treatment outcomes and cost estimate for managing clinical measles in children admitted at Mulago Hospital: A retrospective cohort study. *PLOS Global Public Health*. 2023 Jul 21;3(7):e0001523.
16. Sharma MK, Bhatia V, Swami HM. Outbreak of measles amongst vaccinated children in a slum of Chandigarh.
17. Satpathy SK, Chakraborty AK. Epidemiological study of measles in Singur, West Bengal. *The Journal of Communicable Diseases*. 1990 Mar 1;22(1):23-6.
18. Aurangzeb B, Nisar YB, Hazir T, Burki F, Hassan M. Clinical outcome in children hospitalized with complicated measles. *J Coll Physicians Surg Pak*. 2005 Sep 1;15(9):547-1.
19. Hirfanoğlu T, Tanır G, Karacan C, Göl N. Clinical characteristics, complications and prognosis of seventy-nine measles cases. *Ankara Üniversitesi Tıp Fakültesi Mecmuası*. 2006;59(3):98-103.
20. Deepa KS. Clinical profile of measles in children admitted to a rural tertiary care hospital. *Journal of Evolution of Medical and Dental Sciences*. 2015 Jun 8;4(46):7995-8000.
21. MacKechnie GP, Dalton M, Delport D, Vaccher S. Using Measles Outbreaks to Identify Under-Resourced Health Systems in Low-and Middle-Income Countries: A Predictive Model. *Vaccines*. 2025 Mar 30;13(4):367.
22. Bishaia D, Koenig M, Khan MA. Measles vaccination improves the equity of health outcomes: evidence from Bangladesh. *Health economics*. 2003 May;12(5):415-9.
23. Black RE, Allen LH, Bhutta ZA, Caulfield LE, De Onis M, Ezzati M, Mathers C, Rivera J. Maternal and child undernutrition: global and regional exposures and health consequences. *The lancet*. 2008 Jan 19;371(9608):243-60.
24. Kassebaum NJ, Jasrasaria R, Naghavi M, Wulf SK, Johns N, Lozano R, Regan M, Weatherall D, Chou DP, Eisele TP, Flaxman SR. A systematic analysis of global anemia burden from 1990 to 2010. *Blood*. 2014 Jan 30;123(5):615-24.
25. Rahat F, Abiduzzaman MF, Choudhury AM. Clinical Characteristics of Measles in Infancy: A Hospital Based Study. *Dhaka Shishu (Children) Hospital Journal*. 2020;36(1):20-4.
26. Hafiz MY, Muhammad A, Nadia A, Farrukh M, Shakeela A. Vaccination status of measles diagnosed children admitted in Sheikh Zayed Hospital Rahim Yar Khan.
27. Rudan I, Boschi-Pinto C, Biloglav Z, Mulholland K, Campbell H. Epidemiology and etiology of childhood pneumonia. *Bulletin of the world health organization*. 2008;86:408-16B.
28. Gershon AA. Measles virus (rubeola). In *Mandell, Douglas, and Bennett's principles and practice of infectious diseases 2015 Jan 1* (pp. 1967-1973). WB Saunders.
29. Moss WJ. Measles. *Lancet*. 2017 Dec 2;390(10111):2490-2502. doi: 10.1016/S0140-6736(17)31463-0.
30. Husada D, Kusdwijono, Puspitasari D, Kartina L, Basuki PS, Ismoedijanto. An evaluation of the clinical features of measles virus infection for diagnosis in children within a limited resources setting. *BMC pediatrics*. 2020 Jan 6;20(1):5.
31. Kimura H, Shirabe K, Takeda M, Kobayashi M, Tsukagoshi H, Okayama K, Ryo A, Nagasawa K, Okabe N, Minagawa H, Kozawa K. The association between documentation of Koplik spots and laboratory diagnosis of measles and other rash diseases in a national measles surveillance program in Japan. *Frontiers in Microbiology*. 2019 Feb 18;10:269.
32. Anis-ur-Rehman ST, Idris M. Clinical outcome in measles patients hospitalized with complications. *J Ayub Med Coll Abbottabad*. 2008; 20(2):14-6.